

Wall-associated Kinase WAK1 Interacts with Cell Wall Pectins in a Calcium-induced Conformation

Annabelle Decreux and Johan Messiaen¹

FUNDP, Unité de Recherche en Biologie Végétale, 61 rue de Bruxelles, 5000 Namur, Belgium

Wall-associated kinase 1 (WAK1) is a transmembrane protein containing a cytoplasmic Ser/Thr kinase domain and an extracellular domain in contact with the pectin fraction of the plant cell walls. In order to characterize further the interaction of WAK1 with pectin, a 564 bp DNA sequence corresponding to amino acids 67–254 of the extracellular domain of WAK1 from *Arabidopsis thaliana* was cloned and expressed as a soluble recombinant peptide in yeast. Using enzyme-linked immunosorbent assays (ELISA), we show that peptide WAK_{67–254} binds to polygalacturonic acid (PGA), oligogalacturonides, pectins extracted from *A. thaliana* cell walls and to structurally related alginates. Our results suggest that both ionic and steric interactions are required to match the relatively linear pectin backbone. Binding of WAK_{67–254} to PGA, oligogalacturonides and alginates occurred only in the presence of calcium and in ionic conditions promoting the formation of calcium bridges between oligo- and polymers (also known as ‘egg-boxes’). The conditions inhibiting the formation of calcium bridges (EDTA treatment, calcium substitution, high NaCl concentrations, depolymerization and methylesterification of pectins) also inhibited the binding of WAK_{67–254} to calcium-induced egg-boxes. The relevance of this non-covalent link between WAK_{67–254} and cell wall pectins is discussed in terms of cell elongation, cell differentiation and host–pathogen interactions.

Keywords: *Arabidopsis thaliana* — Cell wall — Egg-box — Pectin — Receptor kinase — Wall-associated kinase

Abbreviations: DM, degree of methylesterification; DP, degree of polymerization; ELISA, enzyme-linked immunosorbent assay; GRP, glycine-rich protein; Gula, guluronic acid; HRGP, hydroxyproline-rich glycoprotein; INA, 2,6-dichloroisonicotinic acid; ManA, manuronic acid; PGA, polygalacturonic acid; PMSF, phenylmethylsulfonyl fluoride; TMB, 3,3',5,5'-tetramethylbenzidine; WAK, wall-associated kinase; WAKL, WAK-like.

Introduction

Analysis of the *Arabidopsis* genome unravelled several hundred open reading frames encoding putative receptor kinases (Morris and Walker 2003). Except for a limited number

of receptor kinases involved in specific gene-for-gene (Song et al. 1995, Liu et al. 2002) or general pathogen recognition (Gomez-Gomez et al. 2001, Gomez-Gomez and Boller 2002), the perception of brassinosteroids (He et al. 2000, Clouse 2002), cytokinins (Yamada et al. 2001, Schafer and Schmulling 2002) ethylene (Gamble et al. 1998, Hua and Meyerowitz 1998, Hua et al. 1998) and systemin (Scheer and Ryan 2002), and those controlling pollen germination (Giranton et al. 2000, Kemp and Doughty 2003), somatic embryogenesis (Hecht et al. 2001), nodulation (Endre et al. 2002, Krusell et al. 2002, Stracke et al. 2002) and meristem identity (Rojo et al. 2002), most receptor kinases remain doubly orphan. The ligand and the signal transduction pathways they control are largely unknown.

The wall-associated kinases (WAKs) belong to this family of orphan receptors. The first WAKs described in *Arabidopsis thaliana* are encoded by five genes (*WAK1–WAK5*) forming a 30 kb gene cluster on chromosome 1 (He et al. 1996, He et al. 1999). This gene family has been extended recently to include several new members coding for WAK-like proteins (WAKLs). They are also grouped in clusters and are distributed among the five chromosomes of *Arabidopsis* (Verica et al. 2003). WAKs and WAKLs were also found in pea, tobacco and maize using anti-WAK antibodies (He et al. 1996, Gens et al. 2000).

WAK1–5 encode Ser/Thr kinases embedded in the plasma membrane. The intracellular domain of the five isoforms is relatively well conserved (86%) and contains the typical eukaryotic Ser/Thr kinase signature. The extracellular domain is not as well conserved (40–64%) and is unique to each WAK. Depending on the isoform, the extracellular domain contains several subdomains and shares some homology with proteins found in the extracellular matrix of mammalian cells. All isoforms, however, share an epidermal growth factor 2 (EGF2)-like and an EGF-Ca²⁺-like domain near the transmembrane domain. The positions of the cysteine residues believed to be involved in the formation of disulfide bonds are conserved among the five WAKs. Some degeneration is noticed, however, in the spacing of the conserved residues and within the amino acid sequences laying in between (He et al. 1999, Verica and He 2002).

Most WAKs are expressed throughout the plant. *WAK1*, 2, 3 and 5 are expressed in green organs; *WAK1* and 2 are also weakly expressed in flowers and siliques; and *WAK2* is also expressed in roots. *WAK4*, on the contrary, is only expressed in siliques. Antisense experiments showed that *WAK* expression is crucial for host–pathogen interactions (He et al. 1998), cell

¹ Corresponding author: E-mail, johan.messiaen@fundp.ac.be; Fax, +32-81-724382.

elongation, morphogenesis and plant development (Lally et al. 2001, Wagner and Kohorn 2001). Since WAKs are believed to play a crucial role at the cell wall–plasmalemma interface, it was postulated that the specific expression patterns of *WAK* genes at the cellular and tissue level reflect the diversity of interactions between the extracellular domain of WAKs and specific cell wall components (Kohorn 2001).

Among the five WAKs, WAK1 is the most studied receptor kinase. *WAK1* encodes a 68 kDa protein of 595 amino acids and is found in almost all tissues. Expression of *WAK1* can be induced by salicylic acid or by its homologue, 2,6-dichloroisonicotinic acid (INA). WAK1 also belongs to the PR protein family since its expression needs the NPR1/NIM1 positive regulator. The nature of the signalling link between WAK1 and NPR1/NIM1 is unknown, however. The expression of an antisense or a dominant-negative allele of *WAK1* showed that *WAK1* expression is crucial for plant survival in the presence of exogenous INA. Similarly, ectopic expression of *WAK1* protects plant tissue against lethal accumulation of salicylic acid during plant–pathogen interactions (He et al. 1998). *WAK1* is also up-regulated during systemic acquired resistance (Maleck et al. 2000), by methyljasmonate and ethylene (Schenk et al. 2000). Other stresses such as high temperatures or high salt concentrations have no effect on *WAK1* expression, whereas wounding reduces its expression (He et al. 1998). Expression of *WAK1* is also induced by aluminium treatment, leading to a delayed accumulation of the receptor kinase at the periphery of cortex cells within the elongation zone of the root apex. Interestingly, overexpression of *WAK1* in transgenic *Arabidopsis* plants increased aluminium tolerance (Sivaguru et al. 2003).

Two-hybrid and in vitro binding experiments using a recombinant subdomain (amino acids 178–334) of the extracellular region of WAK1 allowed the identification of a secreted glycine-rich protein (AtGRP3) that specifically interacts with WAK1. Moreover, exogenous GRP3 induced the transcription of *WAK1*, *AtGRP3* and *PR1* genes in *Arabidopsis*. It was also shown that the WAK1–GRP3 complex interacts with the intracellular KAPP phosphatase to form a 500 kDa complex (Park et al. 2001), which is similar to the complexes involved in the regulation of meristem identity (Stone et al. 1998, Trotochaud et al. 1999). The putative signalling role of this WAK1–GRP3–KAPP complex is largely unknown. One of the possible targets of this complex could be the chloroplast oxygen-evolving enhancer protein 2 (OEE2) which becomes phosphorylated in a WAK1/GRP3-dependent manner in *Arabidopsis* (Yang et al. 2003). The physiological significance of this interaction is unknown.

WAKs, and WAK1 in particular, are also connected to cell wall components. The exact nature of this link is still unclear. WAK1 remains strongly attached to the insoluble cell wall fraction. It can be released from the extracellular matrix either by a 4% SDS, 50 mM dithiothreitol (DTT) treatment and boiling (He et al. 1996), or by a pectinase treatment (Wagner and Kohorn 2001), suggesting that WAK1 might be covalently




Fig. 1 Western blot of the purified recombinant extracellular subdomain WAK_{67–254} in the presence of PGA. An equal amount (100 ng) of purified recombinant WAK_{67–254} was incubated during 1 h at room temperature in the presence of PGA prepared in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer. After centrifugation, the supernatant was analysed by SDS–PAGE and Western blotting. Recombinant proteins were detected using an anti-Xpress antibody. Lane 1, WAK_{67–254}; lanes 2–5, WAK_{67–254} + 0.3, 0.5, 2 or 10 μg of PGA, respectively.

bound to pectin. The interaction between WAK1 and pectins was confirmed by using anti-WAK1 and anti-pectin JIM5 and JIM7 antibodies. Both antibodies recognized the same 68 kDa protein band in Western blots of cell wall proteins extracted from pectinase-treated cell walls (Wagner and Kohorn 2001).

In order to gain a better insight into the signalling role of WAK1 at the plasmalemma–cell wall interface, we investigated the interaction between a subdomain of the extracellular domain of WAK1 and polygalacturonic acid and oligogalacturonides in vitro. We show in this report, using enzyme-linked immunosorbent assays (ELISAs), that the extracellular domain of WAK1 is bound to a calcium-induced conformation in polygalacturonic acid, oligogalacturonides and pectins extracted from *Arabidopsis* cell walls. This interaction was prevented by methylesterification, calcium chelators and pectin depolymerization.

Results

Recombinant WAK_{67–254} binds to PGA

A 564 bp DNA sequence encoding amino acids 67–254 which is unique to the extracellular domain of WAK1 (NCBI accession no. AJ009696) was cloned. It excludes the region encoding the EGF-like repeats common to all WAKs. This sequence, called WAK_{67–254}, was overexpressed in yeast, purified and used in the in vitro binding assays.

In order to confirm the binding of the recombinant subdomain of WAK_{67–254} to pectin, we used a modified binding test described by Carpin et al. (2001) to demonstrate the interaction of recombinant isoperoxidases with PGA in the extracellular matrix of *Cucurbita pepo*. Briefly, the purified recombinant protein was incubated for 1 h at room temperature in the presence of PGA and then centrifuged to discard pelleting PGA–WAK_{67–254} complexes. The resulting supernatant was analysed by SDS–PAGE and Western blotting using an anti-Xpress primary antibody. Fig. 1 shows that the recombinant WAK_{67–254} subdomain disappeared from the supernatant when incubated in the presence of increasing amounts of PGA. This indicates that the recombinant extracellular subdomain WAK_{67–254} interacts with PGA and becomes pelletable in a con-




Fig. 2 (a) Direct ELISA test comparing the interaction of increasing amounts of recombinant WAK₆₇₋₂₅₄ with PGA and the interaction of increasing amounts of a recombinant β -galactosidase (lacZ) with PGA prepared in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer (20 μ g of PGA well⁻¹). Results are expressed as a percentage of the response obtained in the presence of PGA. (b) Direct ELISA test showing typical absorbance values obtained with 250 ng of WAK₆₇₋₂₅₄ in the presence of increasing amounts of PGA coated within the microwells.

centration-dependent way. This recombinant subdomain was analysed further in the following experiments.

Binding of the recombinant WAK₆₇₋₂₅₄ to PGA is specific

In order to characterize the interaction of WAK₆₇₋₂₅₄ with PGA, we used a modified direct ELISA binding assay in which PGA was coated onto the wells of a microplate as a polysaccharide trapper. PGA was prepared in ionic conditions (0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer, pH 8.2) promoting the formation of PGA dimers as described by Liners et al. (1989). After a blocking step, the recombinant proteins were added to the wells and incubated during 1 h at room temperature. Recombinant proteins interacting with PGA were detected with the anti-Xpress primary antibody. The recombinant extracellular domain WAK₆₇₋₂₅₄ bound to immobilized PGA, confirming the Western blot result. This interaction was compared with the binding of a recombinant β -galactosidase (lacZ) purified from yeast in the same conditions and containing the same tags as the recombinant WAK₆₇₋₂₅₄. LacZ did not bind to PGA, indicating that the presence of the purification and detection tags on the recombinant WAK₆₇₋₂₅₄ had no effect on its binding to PGA (Fig. 2a, b). Since similar results were obtained with purified and total yeast extracts containing WAK₆₇₋₂₅₄ (not shown), the following binding experiments were performed only in the presence of the purified WAK₆₇₋₂₅₄ peptide.

In a first step, we wanted to know if the binding of WAK₆₇₋₂₅₄ to PGA could be considered as a simple ionic interaction or as a more complex interaction combining ionic and




Fig. 3 Direct ELISA test comparing the interaction of recombinant WAK₆₇₋₂₅₄ (250 ng) with PGA and other polysaccharides prepared in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer (20 μ g of polysaccharide well⁻¹). Results are expressed as a percentage of the response obtained in the presence of PGA.

structural requirements. The ELISA binding assay was thus used to evaluate the interaction of purified WAK₆₇₋₂₅₄ with other uncharged and charged polysaccharides coated into the wells (Fig. 3). WAK₆₇₋₂₅₄ did not bind to gum arabic, a high molecular weight branched and uncharged polysaccharide with a β (1 \rightarrow 3) galactose backbone. WAK₆₇₋₂₅₄ bound partially to dextran sulfate (30% of PGA control), a high molecular weight branched α (1 \rightarrow 6)-D-glucose polysulfated synthetic polysaccharide. We observed that WAK₆₇₋₂₅₄ bound alginates, negatively charged polysaccharides containing β (1 \rightarrow 4)-L-mannuronic acid (ManA) and α (1 \rightarrow 4)-D-guluronic acid (GulA) distributed as blocks of ManA, GulA and heteropolymeric alternating GulA–ManU sequences (Tombs and Harding 1998). The binding of WAK₆₇₋₂₅₄ to alginates was, however, always lower when compared with PGA and was dependent on the GulA content of the tested polysaccharides: WAK₆₇₋₂₅₄ bound alginate A, containing 70% GulA residues, significantly more than alginates B and C containing a maximum of 40% GulA residues.

Since alginates share similar structural, chemical and physical properties with PGA, their interaction with WAK₆₇₋₂₅₄ was investigated further. A competitive ELISA binding assay was used to measure WAK₆₇₋₂₅₄ binding to a competitor polysaccharide in solution. If binding occurs in solution (competition), WAK₆₇₋₂₅₄ would be inhibited by the competitor and would not be available to bind the PGA coating of the microwells. On the contrary, in the absence of competitor binding, WAK₆₇₋₂₅₄ would bind the PGA coating. We observed that the concentration of soluble PGA needed to prevent 50% binding of WAK₆₇₋₂₅₄ (250 ng) to immobilized PGA was about 15 μ g ml⁻¹ (Fig. 4). Here again, alginates behaved differently depend-




Fig. 4 Competition ELISA test comparing the interaction of recombinant WAK₆₇₋₂₅₄ with alginates and PGA. Competitor alginates or PGA were pre-incubated during 1 h at room temperature with WAK₆₇₋₂₅₄ (250 ng) in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer. The competition mix was added to the wells and incubated further within the PGA-coated microwells (20 µg well⁻¹). Results are expressed as a percentage of control without competitor.

ing on their GulA content. Surprisingly, alginate A with a high GulA content, which are known to form strong inter-chain associations between poly(GulA) domains in the presence of calcium (Tombs and Harding 1998), bound to WAK₆₇₋₂₅₄ better than PGA. The binding of WAK₆₇₋₂₅₄ to alginate A in solution was probably overestimated by this competitive ELISA compared with the direct ELISA (Fig. 3) test in which alginates were directly coated onto the microwells. Solutions of alginate A appeared more viscous than PGA and other alginate (B and C) solutions, limiting the mobility of WAK₆₇₋₂₅₄ to the PGA coating. Alginates B and C, which form fewer and more flexible poly(GulA) inter-chain associations in the presence of calcium than alginate A (Tombs and Harding 1998), also bound to WAK₆₇₋₂₅₄. This interaction was, however, only measured in the presence of relatively high concentrations >300 µg ml⁻¹ of alginates B and C.

Taken together, these results show that WAK₆₇₋₂₅₄ bound polyanions with different affinities. The dextran sulfate data also demonstrate that the presence of negative charges on a polysaccharide was not sufficient to explain WAK₆₇₋₂₅₄ binding, and that additional steric constraints certainly account for the specificity of the interaction with PGA. Since both PGA and poly(GulA) blocks in alginates adopt a similar ‘egg-box’ conformation in the presence of calcium (Davis et al. 2003), we investigated the importance of this conformation for the binding of WAK₆₇₋₂₅₄.

Binding of WAK₆₇₋₂₅₄ to pectin is conformation dependent

Since the conformation of PGA in solution depends largely on its ionic environment, we tested the binding of




Fig. 5 Direct ELISA test comparing the interaction of recombinant WAK₆₇₋₂₅₄ (250 ng) with coated PGA prepared in Tris buffers with different Ca²⁺/Na⁺ ratios. Results are expressed as a percentage of the response obtained in the presence of PGA (20 µg well⁻¹) prepared in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer.




Fig. 6 Direct ELISA test comparing the interaction of recombinant WAK₆₇₋₂₅₄ (250 ng) with coated PGA prepared in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer (‘CaCl₂’), in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer + 5 mM EDTA (‘EDTA’) or in 0.5 mM Mg²⁺/150 mM Na⁺ Tris buffer in which Ca²⁺ is replaced by Mg²⁺ (‘MgCl₂’). Results are expressed as a percentage of the response obtained in the presence of PGA (20 µg well⁻¹) prepared in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer.

WAK₆₇₋₂₅₄ to PGA in the ionic conditions used by Liners et al. (1989) to control the formation of inter-chain bridges in the presence of calcium. We observed that WAK₆₇₋₂₅₄ bound to PGA in the 0.5 mM Ca²⁺/50 mM Na⁺ and 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffers, but not in the 0.5 mM Ca²⁺/300 mM Na⁺ Tris buffer (Fig. 5). These results suggest that WAK₆₇₋₂₅₄ bound to PGA only in conditions promoting the formation of calcium intermolecular bridges previously described as the egg-box model (Morris et al. 1982, Powell et al. 1982, Liners et al. 1989).

In order to confirm this hypothesis, we measured the binding of WAK₆₇₋₂₅₄ to PGA in the 0.5 mM Ca²⁺/150 mM




Fig. 7 Competition ELISA test comparing the interaction of recombinant WAK₆₇₋₂₅₄ (250 ng) with methylesterified pectins or NaOH-treated methylesterified pectins (50 $\mu\text{g ml}^{-1}$) before the addition of WAK₆₇₋₂₅₄. Competitor pectins with a degree of methylesterification of 34 (DM34), 65 (DM65) and 90% (DM90) were pre-incubated during 1 h at room temperature with WAK₆₇₋₂₅₄ in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer. The competition mix was incubated further in the presence of coated PGA (20 $\mu\text{g well}^{-1}$). Pre-incubation with PGA (50 $\mu\text{g ml}^{-1}$) was used as a positive competition control. Results are expressed as a percentage of control without competitor.

Na⁺ buffer containing 5 mM EDTA or in the 0.5 mM Mg²⁺/150 mM Na⁺ buffer in which calcium Ca²⁺ was replaced by Mg²⁺. Fig. 6 shows that both treatments suppressed the binding of WAK₆₇₋₂₅₄ to PGA. As previously shown by Liners et al. (1989), EDTA chelates calcium ions and prevents the formation of calcium intermolecular bridges between PGA polymers. In these conditions, the binding of WAK₆₇₋₂₅₄ to isolated PGA polymers was also inhibited. Similarly, hydrated magnesium ions are larger than hydrated calcium ions and are unable to stabilize PGA dimers or multimers in solution (Liners et al. 1992). Therefore, they did not enhance the binding of WAK₆₇₋₂₅₄ to PGA.

A second indication supporting the hypothesis that binding of WAK₆₇₋₂₅₄ occurs on calcium-bridged PGA came from binding tests performed in the presence of methylesterified pectins. Methylesterification masks the carboxyl groups of PGA and therefore interferes with the formation of inter-chain calcium bridges as previously shown by Liners et al. (1992). We showed that pectins with a degree of methylesterification (DM) of 90 or 65% were not binding substrates for WAK₆₇₋₂₅₄ (Fig. 7). Some WAK₆₇₋₂₅₄ binding was observed, however, in the presence of pectins with a DM of 34%, which corresponds almost to the minimum DM compatible with the formation of stable inter-chain calcium bridges as shown by Liners et al. (1992). Moreover, chemical de-esterification of all tested methylesterified pectins before the incubation step with WAK₆₇₋₂₅₄ restored their ability to adopt a calcium-induced conformation and to bind WAK₆₇₋₂₅₄ again.

A third indication that WAK₆₇₋₂₅₄ binds a PGA conformation came from experiments performed in the presence of




Fig. 8 Competition ELISA test comparing the interaction of recombinant WAK₆₇₋₂₅₄ (250 ng) with oligogalacturonides with a degree of polymerization (DP) ≥ 9 or < 9 . Competitor oligogalacturonides were pre-incubated during 1 h at room temperature with WAK₆₇₋₂₅₄ in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer. The competition mix was incubated further in the presence of coated PGA (20 $\mu\text{g well}^{-1}$). Results are expressed as a percentage of control without competitor.

oligogalacturonides which can adopt a stable calcium-induced conformation if their degree of polymerization (DP) is ≥ 9 (Liners et al. 1992). As shown in Fig. 8, WAK₆₇₋₂₅₄ bound only to oligogalacturonides with a DP ≥ 9 . The concentration of oligogalacturonides with a DP ≥ 9 needed to inhibit 50% of the binding of WAK₆₇₋₂₅₄ (250 ng) to PGA was 20 $\mu\text{g ml}^{-1}$. At higher concentrations, oligogalacturonides with a DP ≥ 9 were, however, unable to compete as efficiently as PGA for the recruitment of WAK₆₇₋₂₅₄. This difference in binding properties of WAK₆₇₋₂₅₄ to oligogalacturonides, compared with PGA, probably relies on the size of the minimum binding site recognized by WAK₆₇₋₂₅₄ on its target. The complex formed between WAK₆₇₋₂₅₄ and a single long polymer (PGA) is likely to be more stable than in the presence of a mixture of oligogalacturonides with a DP ≥ 9 . These results suggest again that binding of WAK₆₇₋₂₅₄ occurs on calcium-bridged oligo- or polygalacturonides since oligomers with a DP ≥ 9 are able, unlike oligomers with a DP < 9 , to undergo a calcium-induced change of conformation in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer.

WAK₆₇₋₂₅₄ binds to pectins purified from *Arabidopsis thaliana* cell walls

A competitive ELISA binding assay was used to measure the binding of WAK₆₇₋₂₅₄ to four pectin fractions extracted from *A. thaliana* purified cell walls. Pectins were extracted successively in the presence of water (fraction F₁), oxalate (fraction F₂), HCl (fraction F₃) and NaOH (fraction F₄). Equal concentrations (15 μg of uronic acids ml^{-1}) of each pectin fraction (native or deesterified) were then incubated with WAK₆₇₋₂₅₄ to evaluate their competitor activity (Fig. 9a). We found that WAK₆₇₋₂₅₄ had little binding activity on native




Fig. 9 (a) Competition ELISA test comparing the interaction of recombinant WAK₆₇₋₂₅₄ (250 ng) with *A. thaliana* pectins extracted from cell walls in water (F₁), in ammonium oxalate (F₂), in HCl (F₃) and in NaOH (F₄). Competitor pectin fractions (native or de-esterified, 15 µg ml⁻¹) were pre-incubated during 1 h at room temperature with WAK₆₇₋₂₅₄ in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer. The competition mix was incubated further in the presence of coated PGA (20 µg well⁻¹). (b) Competitive ELISA test of fractions F₁ to F₄ in the presence of the 2F4 monoclonal antibody. Competitor pectin fractions (native or de-esterified, 15 µg ml⁻¹) were pre-incubated during 1 h at room temperature with the antibody in 0.5 mM Ca²⁺/150 mM Na⁺ Tris buffer. The competition mix was incubated further in the presence of coated PGA (20 µg well⁻¹). Pre-incubation with PGA (20 µg ml⁻¹) was used as a positive competition control. Results are expressed as a percentage of control without competitor.

Arabidopsis pectins unless treated with 0.05 M NaOH for de-esterification (except fraction F₄ that had been already extracted in the presence of 0.05 M NaOH), suggesting that *Arabidopsis* pectins are highly methylesterified. The absence of WAK₆₇₋₂₅₄ binding to the oxalate-extractable pectins of fraction F₂ was surprising since oxalate treatment is known to extract calcium-bound pectins from the cell wall (Schols et al. 1995). Fractions F₁–F₄ were therefore analysed further with the 2F4 monoclonal antibody that specifically recognizes the calcium-induced egg-box conformation in pectin (Liners et al. 1989). The competitive ELISA performed in the presence of the 2F4 antibody showed that native *Arabidopsis* pectins of

fraction F₁ are partially methylesterified, contain the egg-box epitope (70% inhibition, Fig. 9b) but were unable to bind WAK₆₇₋₂₅₄ unless deesterified (Fig. 9a). This indicates that the binding of WAK₆₇₋₂₅₄ to pectin is more severely affected by methylesterification than the binding of the 2F4 antibody to its epitope (Liners et al. 1992). Interestingly, F₂ pectins inhibited the 2F4 response only by 50% and this inhibition was not affected by chemical de-esterification. These results indicate that egg-box epitopes in F₂ pectins are either less abundant (not all galacturonic acid residues are involved in egg-box formation) or less accessible to the 2F4 antibody than in pectins from the F₁, F₃ and F₄ fractions. This lower competitor activity of fraction F₂ in the 2F4 antibody assay probably explains the unsuccessful detection of the binding of WAK₆₇₋₂₅₄ to fraction F₂ (Fig. 9a). We also confirmed that native F₃ pectins are highly methylesterified (no competition), but do adopt an egg-box epitope after de-esterification and then become competitors of the 2F4 response (Fig. 9b). This response was consistent with the binding of WAK₆₇₋₂₅₄ to fraction F₃ (Fig. 9a). As expected from the WAK₆₇₋₂₅₄ binding ELISA test, pectins in fraction F₄ strongly inhibited the 2F4 response (Fig. 9b). These competitive ELISA tests confirm that WAK₆₇₋₂₅₄ binding on *Arabidopsis* pectins (i) is severely affected by methylesterification and (ii) occurs on a pectin domain that is very similar to the egg-box epitope recognized by the 2F4 antibody.

Discussion

The central signalling role of the plant cell extracellular matrix has already been suggested in many physiological responses such as plant cell growth, differentiation and defence against invading pathogens (Brownlee 2002). In animal cells, several plasma membrane proteins and integrins are involved in establishing a signalling continuum between the extracellular matrix, the cytosol and the cytoskeleton that coordinates cell differentiation and responses (Pozzi and Zent 2003). Despite many direct or indirect experiments designed to identify molecular or immunological homologues in plants, it became obvious that plant cells possess unique molecular mechanisms to communicate with its unique cell wall polysaccharides and proteins. The interaction of those cell wall components with the interior of the cell through specific transmembrane ‘receptors’ therefore remains largely unknown.

In this study, we focused on the interaction of pectin with the wall-associated Ser/Thr receptor kinase WAK1 involved in cell elongation, morphogenesis (Lally et al. 2001, Wagner and Kohorn 2001) and protection against pathogens (He et al. 1998). The link between WAK1 and the pectin network was first suggested by He et al. (1996) and Wagner and Kohorn (2001). They showed that only relatively harsh extraction methods or pectinase treatments were able to release WAK1 from purified cell walls, leading to the conclusion that pectins were covalently bound to WAK1. Whatever the exact nature of the link between WAK1 and pectin, it might be relevant for the

molecular dialogue at the cell wall–plasmalemma interface. Pectins are indeed major signalling components of the cell wall involved in a series of physiological responses including morphogenesis, cell differentiation and host–pathogen interactions (Ridley et al. 2001). We showed in this study that PGA binds to a subdomain of the extracellular domain of WAK1 that is different from the binding domain of AtGRP3 (Park et al. 2001). Interestingly, WAK_{67–254} did not bind 65 or 90% methylesterified pectins, suggesting that the negative charges of the polyanion are somehow involved in the assembly of the PGA–WAK_{67–254} complex. This behaviour was also observed with zucchini cell wall isoperoxidases that bind PGA, but not methylesterified pectins (Penel and Greppin 1994). Binding of WAK_{67–254} to PGA was, however, not strictly dependent on the presence of negative charges since WAK_{67–254} did not bind other polyanions *in vitro* with the same affinity.

Binding of WAK_{67–254} to PGA was modulated by calcium ions and, more accurately, by the calcium/monovalent cation ratio. Binding could only be observed in ionic conditions promoting the formation of calcium-bridged PGA dimers and/or multimers (also known as ‘egg-boxes’). Increasing the monovalent cation content, which inhibits the formation of intermolecular bridges between PGA polymers (Liners et al. 1989), also inhibits the interaction with WAK_{67–254}. On the other hand, decreasing the monovalent cation content, which induces the formation of PGA multimers (Liners et al. 1989), reduced by 25% the binding of WAK_{67–254} to PGA. Total removal of calcium ions from the binding environment, replacement of Ca²⁺ ions by Mg²⁺ ions and methylesterification of pectin abolished the formation of PGA–WAK_{67–254} complexes. This calcium-dependent interaction indicates that the binding sites at the surface of the extracellular domain of WAK1 do not simply match negative charges aligned on a single PGA polymer but match a conformation located on two or more PGA chains held together by calcium. The differential binding of WAK_{67–254} to alginates is therefore particularly relevant since poly(GulA) blocks in alginates from brown seaweeds are almost the mirror image of PGA in higher plants. The probability that poly(GulA) blocks form egg-box junction zones in the presence of calcium, similar to those found in PGA, and recruit WAK_{67–254} on this conformation, is therefore higher in alginates containing 70% GulA compared with alginates containing a maximum of 40% GulA. Even if WAK1 has not been identified in brown seaweeds, it is interesting to note here that several secreted proteins from asymmetrically growing *Fucus distichus* embryos share some homology with cell surface proteins, including WAKs, involved in cell morphogenesis in other eukaryotes (Belanger et al. 2003).

Even if we cannot exclude a direct influence of the size of the ligand on WAK_{67–254} recruitment, the results obtained in the presence of oligogalacturonides with a DP ≥ 9 also support the hypothesis that WAK1 interacts with calcium-bridged PGA since WAK_{67–254} bound only to pectin oligomers known to adopt a calcium-induced conformation as described by Liners

et al. (1992). Finally, the binding of WAK_{67–254} to *Arabidopsis* pectins could be clearly correlated with the ability of those pectins to be detected by the 2F4 monoclonal antibody which specifically recognized the egg-box conformation in pectin (Liners et al. 1992).

Our results thus suggest that binding of WAK1 occurs on a calcium-induced conformation of pectin and could be more labile than first mentioned by He et al. (1996). Our data are not incompatible with the results of the above-mentioned study, since the difficulty in extracting WAK1 from the cell wall could be attributed to the use of a 500 mM CaCl₂-containing buffer to resuspend purified cell walls. Considering the calcium-dependent gelling properties of pectin and the importance of calcium for the formation of WAK1–PGA complexes, these ionic conditions certainly stiffened the cell wall, therefore limiting WAK1 mobility and extractability. The existence of a non-covalent link between PGA and WAK1 might also be an explanation for the unsuccessful attempts to immunoprecipitate this complex with either anti-WAK or anti-pectin antibodies as reported by Wagner and Kohorn (2001). WAK1 binding to PGA could resemble the relatively strong interaction of some HRGPs (MacDougall et al. 2001) and isoperoxidases in zucchini and *Arabidopsis* (Carpin et al. 2001, Shah et al. 2004) with pectin through cationic amino acids. In the case of the zucchini isoperoxidase, homology modelling and site-directed mutagenesis allowed the identification of three clustered arginines, located away from the active site of the enzyme, directly involved in binding to calcium-bridged PGA (Carpin et al. 2001).

A non-covalent link between WAK1 and pectins is also not necessarily in contradiction to the extraction of WAK1 by pectinase treatments reported by Wagner and Kohorn (2001). As pectinase treatments simply depolymerize pectins in the cell wall, WAK1 could become extractable when the pectic ligand becomes too short to be bridged by calcium ions (DP <9). Subtle changes of cell wall architecture by pectinases during cell wall turnover, or more drastic changes during pathogen attack, could be sensed by WAK1 and become relevant in terms of cell signalling at the cell wall–plasmalemma interface. Moreover, the interaction of WAK1 with oligogalacturonides with a DP ≥ 9 could modulate the many physiological responses induced by pectin elicitors such as ion fluxes (Mathieu et al. 1991, Messiaen et al. 1993, Messiaen and Van Cutsem 1994, Lecourieux et al. 2002, Navazio et al. 2002), oxidative burst (Legendre et al. 1993), accumulation of methyljasmonate (Doares et al. 1995), activation of mitogen-activated protein kinases (Cazale et al. 1999, Droillard et al. 2000), protein phosphorylation (Droillard et al. 1997), transcription of several defence genes (Messiaen and Van Cutsem 1993), auxin-dependent morphogenesis (Bellincampi et al. 2000, Mauro et al. 2002, Spiro et al. 2002) and fruit ripening (Melotto et al. 1994).

The interaction of the pectic polyanion with cell wall- or plasmalemma-located basic proteins might also be at the origin of signalling processes. Circular dichroism studies demon-

strated that basic domains of carrot extensin and basic synthetic peptides were able to cross-link pectin polymers and induce conformational changes that affect their gelling and swelling behaviour in the presence of calcium. It was shown that both electrostatic and more specific interactions determined by the amino acid sequence of the peptides contribute to the observed new rheological properties (MacDougall et al. 2001). Since the subdomain of the extracellular domain of WAK1 used in this study also contains short basic amino acid sequences, the binding of WAK1 to pectins in the presence of calcium could result in in muro disturbances of the pectin network which in turn could generate signals within the cell wall.

Even if we do not know how WAK1 signals, binding to pectins or pectin oligomers might be crucial for the signalling role of the WAK1–GRP3–KAPP complex described in several physiological responses (Wagner and Kohorn 2001). Although it has been shown that oligogalacturonides bind to cell walls (Mathieu et al. 1998) and plasma membranes (Reymond et al. 1995), and are internalized in a saturable and temperature-dependent way by protoplasts (Horn et al. 1989), no direct link has been made between this binding, a putative oligogalacturonide receptor and a biological response induced by pectin oligomers. Considering the pleiotropic effects of oligogalacturonides on plant cells, WAK1 might belong to one of the perception mechanisms used by the protoplast to sense directly or indirectly the pectins deposited in the cell wall and to respond efficiently to environmental cues that alter pectin content, structure or biological activity. Enzymatic depolymerization of pectins, deposition of newly synthesized methylesterified pectins, de-esterification of pectins in muro, local or systemic changes of the cell wall ionic environment (e.g. through the opening of ion channels), proton pumping or abiotic stresses (heavy metals, high salt concentrations) could modulate the reported signalling role of WAK1 during cell differentiation (Lally et al. 2001, Wagner and Kohorn 2001) and plant–pathogen interactions (He et al. 1998).

Materials and Methods

Polysaccharide and oligogalacturonide preparation

Methylesterified pectin (DM 34, 65 and 90%) and PGA were from Sigma (U.S.A.). Alginate A from *Laminaria hyperborea* with 70% guluronic acid residues was from Sigma (U.S.A.). Alginates B and C from *Macrocystis pyrifera* with maximum 40% GulA residues were from Sigma (U.S.A.) and Fluka (Belgium), respectively.

Methylesterified pectins were de-esterified in the presence of 50 mM NaOH (pH 12) during 30 min at 4°C as described by Liners et al. (1992).

Oligogalacturonides with DP ≥ 9 and DP < 9 were obtained by acid hydrolysis of PGA and HPLC purification as previously described by Messiaen and Van Cutsem (1994).

Oligo- and polysaccharides were prepared in Tris buffer (20 mM Tris–HCl, pH 8.2) supplemented with the appropriate mono- and divalent cations to be tested.

Cloning of the extracellular domain of WAK1 and transformation of Saccharomyces cerevisiae

Total RNA was isolated from 2-week-old *A. thaliana* seedlings (Col-0, N1092) by using Tripure (Roche Applied Science, Belgium) according to the supplier's instructions. First-strand cDNA was synthesized by using the First-Strand Synthesis Kit (Amersham Biosciences, Belgium) according to the supplier's instructions. A 564 bp DNA sequence encoding amino acids 67–254 from the extracellular domain of WAK1 (NCBI accession no. AJ009696) was amplified from first-strand cDNA by polymerase chain reaction (PCR) using *Taq* polymerase (Roche Applied Science, Belgium) and the following primers: 5'-CCCCTCGAGGAGACAAGTGCTGTCCCA-3' and 5'-CCCGGATCCAGGCCACATGTCTTAAGCGAC-3' (Eurogentec, Belgium). This PCR product was ligated into the multiple cloning site of the yeast expression vector pYES2/NT C (Invitrogen, Belgium) designed to express a recombinant protein in fusion with N- and C-terminal poly(His) tags for purification and with an N-terminal Xpress epitope for immunodetection. *Saccharomyces cerevisiae* host cells (strain INVSc1, Invitrogen, Belgium) were transformed with the resulting plasmid using the lithium acetate transformation method according to the supplier's instructions. Recombinant yeast cells were maintained on SC-U medium (SC minimal medium lacking uracil, Invitrogen, Belgium) containing 2% raffinose.

Expression of recombinant protein

Recombinant yeast cells were pre-cultured in 50 ml of liquid SC-U medium containing 2% raffinose at 30°C with shaking. After 24 h, cells were diluted with the induction medium (SC-U medium containing 2% galactose and 1% raffinose) to obtain an OD₆₀₀ of 0.4. After an additional 17 h culture at 30°C with shaking, induced cells were pelleted at 1,500×g for 5 min at 4°C. The cells were then washed with sterile water and centrifuged at 1,500×g for 5 min at 4°C. Yeast pellets were used directly for recombinant protein extraction or were frozen in liquid nitrogen and stored at –80°C until use. A pYES2/NT/lacZ vector was used as a positive control for protein expression and as negative control for the binding tests.

Total protein extraction in native conditions

Total proteins were extracted by grinding the cells in liquid nitrogen in the presence of 3.5 ml of non-denaturing lysis buffer [140 mM NaCl, 2.7 mM KCl, 20 mM Tris–HCl (pH 7.9), 1 mM phenylmethylsulfonyl fluoride (PMSF), 1 µg ml⁻¹ aprotinin, 1 µg ml⁻¹ chymostatin, 1 µM pepstatin A, 100 µM leupeptin]. The cell lysate was centrifuged at 3,000×g for 15 min at 4°C and the resulting supernatant was used in the ELISA tests.

Purification of recombinant protein

Total proteins were extracted by grinding the cells in liquid nitrogen in the presence of 7 ml of denaturing guanidinium lysis buffer [6 M guanidine-HCl, 20 mM sodium phosphate buffer (pH 7.8), 500 mM NaCl, 1 mM PMSF, 1 µg ml⁻¹ aprotinin, 1 µg ml⁻¹ chymostatin, 1 µM pepstatin A, 100 µM leupeptin]. The resulting lysate was incubated for 10 min at room temperature under mild agitation and then centrifuged at 3,000×g for 15 min at 4°C. The recombinant fusion protein was purified from the supernatant on a nickel-chelating resin according to the supplier's instructions (Probond resine, Invitrogen, Belgium). The eluate containing the recombinant protein was dialysed sequentially overnight as described by Ferguson and Goodrich (2001). The final dialysis was performed in the appropriate buffer used in the ELISA tests. Binding activity of the dialysed and purified WAK_{67–254} peptide to PGA was compared with the binding activity of the non-purified native WAK_{67–254} peptide by ELISA before use in further experiments (positive renaturation control). Protein concentration was

determined with the NanoOrange protein quantification kit (Molecular Probes, U.S.A.).

Direct ELISA binding assay

Maxisorp microplates (VWR, Belgium) were pre-treated with polylysine-HBr (50 $\mu\text{g ml}^{-1}$ in H_2O , 50 $\mu\text{l well}^{-1}$, Sigma, U.S.A.) during 1 h at room temperature. The wells were washed once with 250 $\mu\text{l well}^{-1}$ of 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer (20 mM Tris–HCl, 150 mM NaCl, 0.5 mM CaCl_2 , pH 8.2) and coated overnight at 4°C with the trapper polysaccharide at the appropriate concentration in 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer (50 $\mu\text{l well}^{-1}$) as described by Liners et al. (1989). Non-specific binding sites were blocked for 2 h at room temperature with 3% low fat dried milk dissolved in the 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer. After removal of excess blocking solution, the recombinant proteins were added and incubated for 2 h at room temperature. The wells were washed with 5×250 μl of 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer and further incubated during 1 h at room temperature with 1 $\mu\text{g ml}^{-1}$ of anti-Xpress primary antibody (Invitrogen, Belgium) in the 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer containing 1% low fat dried milk (50 $\mu\text{l well}^{-1}$). After washing with 7×250 μl of 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer, 50 μl of a 1/1,000 dilution of horseradish peroxidase (HRP)-SAM secondary antibody (Amersham Biosciences, Belgium) prepared in 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer containing 1% low fat dried milk were added and incubated for 1 h at room temperature. After washing with 7×250 μl of 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer, the binding of the recombinant protein to the coated polysaccharide was visualized in the presence of the TMB substrate (Enhanced K Blue TMB substrate, Neogen, U.S.A.). Absorbance was measured after 20 min in the dark at 650 nm with a microplate reader (Elx800, Bio-TEK instruments, U.S.A.). Binding assays were done in triplicate for at least three protein extracts.

Competitive ELISA binding assay

When compared with the binding assay using direct ELISA, interaction between the recombinant protein and the polysaccharide occurs in solution. Excess or non-interacting recombinant proteins are detected by their binding on the coated trapper polysaccharide. Recombinant proteins were therefore pre-incubated for 1 h with different competitor oligo- or polysaccharides prepared at the appropriate dilution in 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer. Plates were coated with PGA and blocked with 3% low fat dried milk dissolved in the 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer as described by Liners et al. (1989). The competition mixtures were then added to the wells and left for 2 h at room temperature. The following steps are strictly identical to the direct ELISA binding assay.

Competitive ELISA tests with the 2F4 monoclonal antibody were performed as described by Liners et al. (1989).

Western blot

Purified recombinant proteins were pre-incubated for 1 h at room temperature with increasing amounts of PGA prepared in 0.5 mM Ca^{2+} /150 mM Na^+ Tris buffer (pH 8.2). After centrifugation at 8,500×g, the supernatants were analyzed by SDS–PAGE according to Laemmli (1970) on 12% polyacrylamide gels and electroblotted on a Hybond-P transfer membrane (Amersham Biosciences, Belgium) using a trans-blot semi-dry electrophoretic transfer cell (Biorad, Belgium). After blocking with 3% low fat dried milk prepared in TBS buffer (25 mM Tris, 137 mM NaCl, 2.7 mM KCl, pH 7.4), the membrane was incubated for 1 h with 0.2 $\mu\text{g ml}^{-1}$ anti-Xpress primary antibody (Invitrogen, Belgium) in TBS containing 1% low fat dried milk. The blot was washed three times with TBS and incubated for 1 h with HRP–SAM secondary antibody (Amersham Biosciences, Belgium) diluted 1/5,000 in TBS 1% low fat dried milk. Membranes were

washed three times in TBS buffer and developed with the ECL Plus system (Amersham Biosciences, Belgium), and autoradiographed with BioMax Films (Kodak).

Pectin extraction from *Arabidopsis thaliana* cell walls

Ten-day-old *A. thaliana* (ecotype A-TL) suspension-cultured cells (20 g fresh weight) were ground in 100 ml absolute ethanol and transferred in 100 ml of boiling absolute ethanol during 20 min. After centrifugation (3,000×g, 5 min), the pellet was washed twice with 100 ml of aqueous ethanol (70%; v/v) and incubated overnight at 4°C with 100 ml of aqueous ethanol (70%; v/v). The insoluble cell wall material was then washed with absolute ethanol and acetone and allowed to dry at room temperature. Pectins were extracted successively from 300 mg of dried cell wall material in the presence of distilled water (fraction F_1) for 0.5 h at room temperature, 1% ammonium oxalate (fraction F_2) for 0.5 h at room temperature, 0.05 M HCl for 0.5 h at 85°C (fraction F_3) and 0.05 M NaOH for 0.5 h at 4°C (fraction F_4). Pectin fractions were neutralized, if necessary, as described by Rombouts and Thibault (1986). The uronic acid content of each fraction was determined with the meta-hydroxybiphenyl method according to Blumenkrantz and Asboe-Hansen (1973).

Acknowledgments

We wish to thank Professor P. Van Cutsem and Dr. F. Liners for their advices, valuable comments and for generously providing the 2F4 monoclonal antibody. We also thank C. Demazy and P. Cambier for technical assistance. This work was supported by the Fond National de la Recherche Scientifique (FRFC 2.4.501.00.F)

References

- Belanger, K.D., Wyman, A.J., Sudol, M.N., Singla-Pareek, S.L. and Quatrano, R.S. (2003) A signal peptide secretion screen in *Fucus distichus* embryos reveals expression of glucanase, EGF domain-containing, and LRR receptor kinase-like polypeptides during asymmetric cell growth. *Planta* 217: 931–950.
- Bellincampi, D., Dipierro, N., Salvi, G., Cervone, F. and De Lorenzo, G. (2000) Extracellular H_2O_2 induced by oligogalacturonides is not involved in the inhibition of the auxin-regulated roIB gene expression in tobacco leaf explants. *Plant Physiol.* 122: 1379–1385.
- Blumenkrantz, N. and Asboe-Hansen, G. (1973) New method for quantitative determination of uronic acids. *Anal. Biochem.* 54: 484–489.
- Brownlee, C. (2002) Role of the extracellular matrix in cell-cell signalling: paracrine paradigms. *Curr. Opin. Plant Biol.* 5: 396–401.
- Carpin, S., Crevecoeur, M., de Meyer, M., Simon, P., Greppin, H. and Penel, C. (2001) Identification of a Ca^{2+} -pectate binding site on an apoplastic peroxidase. *Plant Cell* 13: 511–520.
- Cazale, A.C., Droillard, M.J., Wilson, C., Heberle-Bors, E., Barbier-Brygoo, H. and Lauriere, C. (1999) MAP kinase activation by hypoosmotic stress of tobacco cell suspensions: towards the oxidative burst response? *Plant J.* 19: 297–307.
- Clouse, S.D. (2002) Brassinosteroid signal transduction: clarifying the pathway from ligand perception to gene expression. *Mol. Cell* 10: 973–982.
- Davis, T.A., Volesky, B. and Mucci, A. (2003) A review of the biochemistry of heavy metal biosorption by brown algae. *Water Res.* 37: 4311–4330
- Doares, S.H., Syrovets, T., Weiler, E.W. and Ryan, C.A. (1995) Oligogalacturonides and chitosan activate plant defensive genes through the octadecanoid pathway. *Proc. Natl Acad. Sci. USA* 92: 4095–4098.
- Droillard, M.J., Guclu, J., Le Caer, J.P., Mathieu, Y., Guern, J. and Lauriere, C. (1997) Identification of calreticulin-like protein as one of the phosphoproteins modulated in response to oligogalacturonides in tobacco cells. *Planta* 202: 341–348.
- Droillard, M.J., Thibivilliers, S., Cazale, A.C., Barbier-Brygoo, H. and Lauriere, C. (2000) Protein kinases induced by osmotic stresses and elicitor mole-

- cules in tobacco cell suspensions: two crossroad MAP kinases and one osmoregulation-specific protein kinase. *FEBS Lett.* 474: 217–222.
- Endre, G., Kereszt, A., Kevei, Z., Mihacea, S., Kalo, P. and Kiss, G.B. (2002) A receptor kinase gene regulating symbiotic nodule development. *Nature* 417: 962–966.
- Ferguson, H.A. and Goodrich, J.A. (2001) Expression and purification of recombinant human c-Fos/c-Jun that is highly active in DNA binding and transcriptional activation in vivo. *Nucl. Acids Res.* 29: e98
- Gamble, R.L., Coonfield, M.L. and Schaller, G.E. (1998) Histidine kinase activity of the ETR1 ethylene receptor from *Arabidopsis*. *Proc. Natl Acad. Sci. USA* 95: 7825–7829.
- Gens, J.S., Fujiki, M. and Pickard, B.G. (2000) Arabinogalactan protein and wall-associated kinase in a plasmalemmal reticulum with specialized vertices. *Protoplasma* 212: 115–134.
- Giranton, J.L., Dumas, C., Cock, J.M. and Gaude, T. (2000) The integral membrane S-locus receptor kinase of *Brassica* has serine/threonine kinase activity in a membranous environment and spontaneously forms oligomers in planta. *Proc. Natl Acad. Sci. USA* 97: 3759–3764.
- Gomez-Gomez, L., Bauer, Z. and Boller, T. (2001) Both the extracellular leucine-rich repeat domain and the kinase activity of FLS2 are required for flagellin binding and signaling in *Arabidopsis*. *Plant Cell* 13: 1155–1163.
- Gomez-Gomez, L. and Boller, T. (2002) Flagellin perception: a paradigm for innate immunity. *Trends Plant Sci.* 7: 251–256.
- He, Z., Wang, Z.Y., Li, J., Zhu, Q., Lamb, C., Ronald, P. and Chory, J. (2000) Perception of brassinosteroids by the extracellular domain of the receptor kinase BRI1. *Science* 288: 2360–2363.
- He, Z.H., Cheeseman, I., He, D. and Kohorn, B.D. (1999) A cluster of five cell wall-associated receptor kinase genes, Wak1-5, are expressed in specific organs of *Arabidopsis*. *Plant Mol. Biol.* 39: 1189–1196.
- He, Z.H., Fujiki, M. and Kohorn, B.D. (1996) A cell wall-associated, receptor-like protein kinase. *J. Biol. Chem.* 271: 19789–19793.
- He, Z.H., He, D. and Kohorn, B.D. (1998) Requirement for the induced expression of a cell wall associated receptor kinase for survival during the pathogen response. *Plant J.* 14: 55–63.
- Hecht, V., Vielle-Calzada, J.P., Hartog, M.V., Schmidt, E.D., Boutilier, K., Grossniklaus, U. and de Vries, S.C. (2001) The *Arabidopsis* SOMATIC EMBRYOGENESIS RECEPTOR KINASE 1 gene is expressed in developing ovules and embryos and enhances embryogenic competence in culture. *Plant Physiol.* 127: 803–816.
- Horn, M.A., Heinstein, P.F. and Low, P.S. (1989) Receptor-mediated endocytosis in plant cells. *Plant Cell* 1: 1003–1009.
- Hua, J. and Meyerowitz, E.M. (1998) Ethylene responses are negatively regulated by a receptor gene family in *Arabidopsis thaliana*. *Cell* 94: 261–271.
- Hua, J., Sakai, H., Nourizadeh, S., Chen, Q.G., Bleecker, A.B., Ecker, J.R. and Meyerowitz, E.M. (1998) EIN4 and ERS2 are members of the putative ethylene receptor gene family in *Arabidopsis*. *Plant Cell* 10: 1321–1332.
- Kemp, B.P. and Doughty, J. (2003) Just how complex is the *Brassica* S-receptor complex? *J. Exp. Bot.* 54: 157–168.
- Kohorn, B.D. (2001) WAKs; cell wall associated kinases. *Curr. Opin. Cell Biol.* 13: 529–533.
- Krusell, L., Madsen, L.H., Sato, S., Aubert, G., Genua, A., Szczyglowski, K., Duc, G., Kaneko, T., Tabata, S., de Bruijn, F., Pajuelo, E., Sandal, N. and Stougaard, J. (2002) Shoot control of root development and nodulation is mediated by a receptor-like kinase. *Nature* 420: 422–426.
- Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680–685
- Lally, D., Ingmire, P., Tong, H.Y. and He, Z.H. (2001) Antisense expression of a cell wall-associated protein kinase, WAK4, inhibits cell elongation and alters morphology. *Plant Cell* 13: 1317–1331.
- Lecourieux, D., Mazars, C., Pauly, N., Ranjeva, R. and Pugin, A. (2002) Analysis and effects of cytosolic free calcium increases in response to elicitors in *Nicotiana glauca* cells. *Plant Cell* 14: 2627–2641.
- Legendre, L., Rueter, S., Heinstein, P.F. and Low, P.S. (1993) Characterization of the oligogalacturonide-induced oxidative burst in cultured soybean (*Glycine max*) cells. *Plant Physiol.* 102: 233–240.
- Liners, F., Letesson, J.J., Didembourg, C. and Van Cutsem, P. (1989) Monoclonal-antibodies against pectin—recognition of a conformation induced by calcium. *Plant Physiol.* 91: 1419–1424.
- Liners, F., Thibault, J.F. and Van Cutsem, P. (1992) Influence of the degree of polymerization of oligogalacturonates and of esterification pattern of pectin on their recognition by monoclonal antibodies. *Plant Physiol.* 99: 1099–1104.
- Liu, G.Z., Pi, L.Y., Walker, J.C., Ronald, P.C. and Song, W.Y. (2002) Biochemical characterization of the kinase domain of the rice disease resistance receptor-like kinase XA21. *J. Biol. Chem.* 277: 20264–20269.
- MacDougall, A.J., Brett, G.M., Morris, V.J., Rigby, N.M., Ridout, M.J. and Ring, S.G. (2001) The effect of peptide–pectin interactions on the gelation behaviour of a plant cell wall pectin. *Carbohydr. Res.* 335: 115–126.
- Maleck, K., Levine, A., Eulgem, T., Morgan, A., Schmid, J., Lawton, K.A., Dangl, J.L. and Dietrich, R.A. (2000) The transcriptome of *Arabidopsis thaliana* during systemic acquired resistance. *Nat. Genet.* 26: 403–410.
- Mathieu, Y., Guern, J., Spiro, M.D., O'Neill, M.A., Kates, K., Darvill, A.G. and Albersheim, P. (1998) The transient nature of the oligogalacturonide-induced ion fluxes of tobacco cells is not correlated with fragmentation of the oligogalacturonides. *Plant J.* 16: 305–311.
- Mathieu, Y., Kurkdjian, A., Xia, H., Guern, J., Koller, A., Spiro, M., O'Neill, M., Albersheim, P. and Darvill, A. (1991) Membrane responses induced by oligogalacturonides in suspension-cultured tobacco cells. *Plant J.* 1: 333–343.
- Mauro, M.L., De Lorenzo, G., Costantino, P. and Bellincampi, D. (2002) Oligogalacturonides inhibit the induction of late but not of early auxin-responsive genes in tobacco. *Planta* 215: 494–501.
- Melotto, E., Greve, L.C. and Labavitch, J.M. (1994) Cell wall metabolism in ripening fruit (vii. biologically active pectin oligomers in ripening tomato (*Lycopersicon esculentum* Mill.) fruits). *Plant Physiol.* 106: 575–581.
- Messiaen, J., Read, N.D., Van Cutsem, P. and Trewavas, A.J. (1993) Cell wall elicitors increase cytosolic free Ca²⁺ in plant cells. *J. Cell Sci.* 104: 365–371.
- Messiaen, J. and Van Cutsem, P. (1993) Defense gene transcription in carrot cells treated with oligogalacturonides. *Plant Cell Physiol.* 34: 1117–1123.
- Messiaen, J. and Van Cutsem, P. (1994) Pectic signal transduction in carrot cells: membrane, cytosolic and nuclear responses induced by oligogalacturonides. *Plant Cell Physiol.* 35: 677–689.
- Morris, E.R., Powell, D.A., Gidley, M.J. and Rees, D.A. (1982) Conformations and interactions of pectins I. Polymorphism between gel and solid states of calcium polygalacturonate. *J. Mol. Biol.* 155: 507–516.
- Morris, E.R. and Walker, J.C. (2003) Receptor-like protein kinases: the keys to response. *Curr. Opin. Plant Biol.* 6: 339–342.
- Navazio, L., Moscattello, R., Bellincampi, D., Baldan, B., Meggio, F., Brini, M., Bowler, C. and Mariani, P. (2002) The role of calcium in oligogalacturonide-activated signalling in soybean cells. *Planta* 215: 596–605.
- Park, A.R., Cho, S.K., Yun, U.J., Jin, M.Y., Lee, S.H., Sachetto-Martins, G. and Park, O.K. (2001) Interaction of the *Arabidopsis* receptor protein kinase Wak1 with a glycine-rich protein, AtGRP-3. *J. Biol. Chem.* 276: 26688–26693.
- Penel, C. and Greppin, H. (1994) Binding of plant isoperoxidase to pectin in the presence of calcium. *FEBS Lett.* 343: 51–55.
- Powell, D.A., Morris, E.R., Gidley, M.J. and Rees, D.A. (1982) Conformations and interactions of pectins II. Influence of residue sequence on chain association in calcium pectate gels. *J. Mol. Biol.* 155: 517–531.
- Pozzi, A. and Zent, R. (2003) Integrins: sensors of extracellular matrix and modulators of cell function. *Nephron Exp. Nephrol.* 94: e77–e84.
- Reymond, P., Grunberger, S., Paul, K., Muller, M. and Farmer, E.E. (1995) Oligogalacturonide defense signals in plants: large fragments interact with the plasma membrane in vitro. *Proc. Natl Acad. Sci. USA* 92: 4145–4149.
- Ridley, B.L., O'Neill, M.A. and Mohnen, D. (2001) Pectins: structure, biosynthesis, and oligogalacturonide-related signaling. *Phytochemistry* 57: 929–967.
- Rojo, E., Sharma, V.K., Kovaleva, V., Raikhel, N.V. and Fletcher, J.C. (2002) CLV3 is localized to the extracellular space, where it activates the *Arabidopsis* CLAVATA stem cell signaling pathway. *Plant Cell* 14: 969–977.
- Rombouts, F.M. and Thibault, J.-F. (1986) Feruloylated pectic substances from sugar beet pulp. *Carbohydr. Res.* 154: 177–187.
- Schafer, S. and Schumling, T. (2002) The CRK1 receptor-like kinase gene of tobacco is negatively regulated by cytokinin. *Plant Mol. Biol.* 50: 155–166.
- Scheer, J.M. and Ryan, C.A. Jr (2002) The systemin receptor SR160 from *Lycopersicon peruvianum* is a member of the LRR receptor kinase family. *Proc. Natl Acad. Sci. USA* 99: 9585–9590.
- Schenk, P.M., Kazan, K., Wilson, I., Anderson, J.P., Richmond, T., Somerville, S.C. and Manners, J.M. (2000) Coordinated plant defense responses in *Arabidopsis* revealed by microarray analysis. *Proc. Natl Acad. Sci. USA* 97: 11655–11660.

- Schols, H.A., Vierhuis, E., Bakx, E.J. and Voragen, A.G.J. (1995) Different populations of pectic hairy regions occur in apple cell walls. *Carbohydr. Res.* 275: 343–360.
- Shah, K., Penel, C., Gagnon, J. and Dunand, C. (2004) Purification and identification of a Ca²⁺-pectate binding peroxidase from *Arabidopsis* leaves. *Phytochemistry* 65: 307–312.
- Sivaguru, M., Ezaki, B., He, Z.H., Tong, H., Osawa, H., Baluska, F., Volkmann, D. and Matsumoto, H. (2003) Aluminum-induced gene expression and protein localization of a cell wall-associated receptor kinase in *Arabidopsis*. *Plant Physiol.* 132: 2256–2266.
- Song, W.Y., Wang, G.L., Chen, L.L., Kim, H.S., Pi, L.Y., Holsten, T., Gardner, J., Wang, B., Zhai, W.X., Zhu, L.H. et al. (1995) A receptor kinase-like protein encoded by the rice disease resistance gene, Xa21. *Science* 270: 1804–1806.
- Spiro, M.D., Bowers, J.F. and Cosgrove, D.J. (2002) A comparison of oligogalacturonide- and auxin-induced extracellular alkalization and growth responses in roots of intact cucumber seedlings. *Plant Physiol.* 130: 895–903.
- Stone, J.M., Trotochaud, A.E., Walker, J.C. and Clark, S.E. (1998) Control of meristem development by CLAVATA1 receptor kinase and kinase-associated protein phosphatase interactions. *Plant Physiol.* 117: 1217–1225.
- Stracke, S., Kistner, C., Yoshida, S., Mulder, L., Sato, S., Kaneko, T., Tabata, S., Sandal, N., Stougaard, J., Szczygłowski, K. and Parniske, M. (2002) A plant receptor-like kinase required for both bacterial and fungal symbiosis. *Nature* 417: 959–962.
- Tombs, M.P. and Harding, S.E. (1998) Marine polysaccharides. In *An Introduction to Polysaccharide Biotechnology*. Edited by Tombs, M.P. and Harding, S.E. pp. 123–154. Taylor and Francis, London
- Trotochaud, A.E., Hao, T., Wu, G., Yang, Z. and Clark, S.E. (1999) The CLAVATA1 receptor-like kinase requires CLAVATA3 for its assembly into a signaling complex that includes KAPP and a Rho-related protein. *Plant Cell* 11: 393–406.
- Verica, J.A., Chae, L., Tong, H., Ingmire, P. and He, Z.H. (2003) Tissue-specific and developmentally regulated expression of a cluster of tandemly arrayed cell wall-associated kinase-like kinase genes in *Arabidopsis*. *Plant Physiol.* 133: 1732–1746.
- Verica, J.A. and He, Z.H. (2002) The cell wall-associated kinase (WAK) and WAK-like kinase gene family. *Plant Physiol.* 129: 455–459.
- Wagner, T.A. and Kohorn, B.D. (2001) Wall-associated kinases are expressed throughout plant development and are required for cell expansion. *Plant Cell* 13: 303–318.
- Yamada, H., Suzuki, T., Terada, K., Takei, K., Ishikawa, K., Miwa, K., Yamashino, T. and Mizuno, T. (2001) The *Arabidopsis* AHK4 histidine kinase is a cytokinin-binding receptor that transduces cytokinin signals across the membrane. *Plant Cell Physiol.* 42: 1017–1023.
- Yang, E.J., Oh, Y.A., Lee, E.S., Park, A.R., Cho, S.K., Yoo, Y.J. and Park, O.K. (2003) Oxygen-evolving enhancer protein 2 is phosphorylated by glycine-rich protein 3/wall-associated kinase 1 in *Arabidopsis*. *Biochem. Biophys. Res. Commun.* 305: 862–868.

(Received March 29, 2004; Accepted October 7, 2004)